

Main panel 1L bottle

---

**CAUTION**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



# Rentokil IGR Grain Protectant

ACTIVE CONSTITUENT: 30g/L S-METHOPRENE  
SOLVENT: 720g/L LIQUID HYDROCARBONS

GROUP	<b>7A</b>	INSECTICIDE
-------	-----------	-------------

For mixing with uninfested cereal grain for protection against immature stages of Lesser Grain Borer, Rust-Red Flour Beetle and Sawtoothed Grain Beetle, including organophosphate-resistant strains.

**Contents 1 Litre**

**RENTOKIL INITIAL Pty Ltd** A.C.N. 000 034 597  
150 Mowbray Road, Willoughby, NSW 2068 Australia  
a Rentokil Initial company  
Telephone 02 9370 9300

**DIRECTIONS FOR USE:**

**Restrains:** DO NOT move treated grain within 24 hours after treatment. Rentokil IGR is only effective against insects in the immature stages, not adults.

Situation/ Crop	Pest	Rate	Critical Comments
Stored Cereal grain	Lesser grain borer, rust-red flour beetle and sawtoothed grain beetle including those resistant to organophosphates.	1L in 50L of water, for 6 to 9 months protection.	Apply at rate of 1L diluted spray per tonne of grain.
	Stored grain insect pests (excluding <i>Sitophilus</i> spp.)	As above plus Rentokil Fenitrothion 1000 insecticide at the rates shown on the label.	

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

**IN WA FOR USE BY BULK HANDLING AUTHORITIES ONLY.**

**WITHHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED**

**GENERAL INSTRUCTIONS**

**Insecticide Resistance Warning**

<b>GROUP</b>	<b>7A</b>	<b>INSECTICIDE</b>
--------------	-----------	--------------------

For insecticide resistance management Rentokil IGR Grain Protectant is a Group 7A insecticide.

Some naturally-occurring insect biotypes resistant to Rentokil IGR Grain Protectant and other Group 7A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Rentokil IGR Grain Protectant or other Group 7A insecticides are used repeatedly. The effectiveness of Rentokil IGR Grain Protectant on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Rentokil Initial Pty Ltd accepts no liability for any losses that may result from the failure of Rentokil IGR Grain Protectant to control resistant insects.

Rentokil IGR Grain Protectant may be subject to specific resistance management strategies. For further information contact your local supplier, Rentokil Initial Pty Ltd representative or local agricultural department agronomist.

**MIXING:**

Using clean water only, dilute sufficient material for the job at hand. Diluted spray should be used within 48 hours.

**EQUIPMENT:**

The diluted spray should be applied through special grain spraying equipment. The spray nozzle(s) must be capable of delivering an even spray pattern, and the output of spray must be regulated according to the flow of the grain.

**CHECK RATE OF TREATMENT:**

The spray rate, measured in litres per hour must equal the auger or elevator uptake in tonnes per hour.

**CLEANING UP:**

On completion, wash out spraying equipment and mixing and measuring utensils with water.

## **EXPORT OF TREATED PRODUCE**

Users should note that suitable MRLs or import tolerances might not have been established in all markets for produce treated with Rentokil IGR. Before treating grain for export, please check with Rentokil Initial Pty Ltd for the latest information on MRLs and import tolerances.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT:** DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

**STORAGE AND DISPOSAL:** Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**SAFETY DIRECTIONS:** Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. Do not inhale vapour or spray mist. When preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length PVC gloves. Wash hands after use. After each day's use, wash gloves and contaminated clothing.

**FIRST AID:** If poisoning occurs, contact a Doctor or Poisons Information Centre. Phone Australia 131126. If swallowed do NOT induce vomiting. Give a glass of water.

**MSDS:** Additional information is listed in the Material Safety Data Sheet.

**MANUFACTURERS WARRANTY AND EXCLUSION OF LIABILITY:** This product as supplied is of high grade and believed to be suitable for any purpose for which it is expressly recommended and must be used in accordance with the Directions for Use given on the label. No responsibility is accepted in respect of the product save those non-excludable conditions implied by any Federal and State legislation or law of a Territory.

**APVMA Approval No: 53038/1/0804**

**Batch No:**

**DOM**

**Bar Code**

*Main panel 10 L bottle*

---

**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

# **Rentokil**

# **IGR**

## **Grain Protectant**

**ACTIVE CONSTITUENT: 30g/L S-METHOPRENE**  
**SOLVENT: 720g/L LIQUID HYDROCARBONS**

<b>GROUP</b>	<b>7A</b>	<b>INSECTICIDE</b>
--------------	-----------	--------------------

For mixing with uninfested cereal grain for protection against immature stages of Lesser Grain Borer, Rust-Red Flour Beetle and Sawtoothed Grain Beetle, including organophosphate-resistant strains.

**Contents 10 litres**

**RENTOKIL INITIAL Pty Ltd** A.C.N. 000 034 597  
150 Mowbray Road, Willoughby, NSW 2068 Australia  
a Rentokil Initial company  
Telephone 02 9370 9300

**DIRECTIONS FOR USE:**

**Restrains:** DO NOT move treated grain within 24 hours after treatment. Rentokil IGR is only effective against insects in the immature stages, not adults.

Situation/ Crop	Pest	Rate	Critical Comments
Stored Cereal grain	Lesser grain borer, rust-red flour beetle and sawtoothed grain beetle including those resistant to organophosphates.	1L in 50L of water, for 6 to 9 months protection.	Apply at rate of 1L diluted spray per tonne of grain.
	Stored grain insect pests (excluding <i>Sitophilus</i> spp.)	As above plus Rentokil Fenitrothion 1000 insecticide at the rates shown on the label.	

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

**IN WA FOR USE BY BULK HANDLING AUTHORITIES ONLY.**

**WITHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED**

**GENERAL INSTRUCTIONS**

**Insecticide Resistance Warning**

<b>GROUP</b>	<b>7A</b>	<b>INSECTICIDE</b>
--------------	-----------	--------------------

For insecticide resistance management Rentokil IGR Grain Protectant is a Group 7A insecticide.

Some naturally-occurring insect biotypes resistant to Rentokil IGR Grain Protectant and other Group 7A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Rentokil IGR Grain Protectant or other Group 7A insecticides are used repeatedly. The effectiveness of Rentokil IGR Grain Protectant on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Rentokil Initial Pty Ltd accepts no liability for any losses that may result from the failure of Rentokil IGR Grain Protectant to control resistant insects.

Rentokil IGR Grain Protectant may be subject to specific resistance management strategies. For further information contact your local supplier, Rentokil Initial Pty Ltd representative or local agricultural department agronomist.

**MIXING:**

Using clean water only, dilute sufficient material for the job at hand. Diluted spray should be used within 48 hours.

**EQUIPMENT:**

The diluted spray should be applied through special grain spraying equipment. The spray nozzle(s) must be capable of delivering an even spray pattern, and the output of spray must be regulated according to the flow of the grain.

**CHECK RATE OF TREATMENT:**

The spray rate, measured in litres per hour must equal the auger or elevator uptake in tonnes per hour.

**CLEANING UP:**

On completion, wash out spraying equipment and mixing and measuring utensils with water.

## **EXPORT OF TREATED PRODUCE**

Users should note that suitable MRLs or import tolerances might not have been established in all markets for produce treated with Rentokil IGR. Before treating grain for export, please check with Rentokil Initial Pty Ltd for the latest information on MRLs and import tolerances.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT:** DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

**STORAGE AND DISPOSAL:** Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**SAFETY DIRECTIONS:** Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. Do not inhale vapour or spray mist. When preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length PVC gloves. Wash hands after use. After each day's use, wash gloves and contaminated clothing.

**FIRST AID:** If poisoning occurs, contact a Doctor or Poisons Information Centre. Phone Australia 131126. If swallowed do NOT induce vomiting. Give a glass of water.

**MSDS:** Additional information is listed in the Material Safety Data Sheet.

**MANUFACTURERS WARRANTY AND EXCLUSION OF LIABILITY:** This product as supplied is of high grade and believed to be suitable for any purpose for which it is expressly recommended and must be used in accordance with the Directions for Use given on the label. No responsibility is accepted in respect of the product save those non-excludable conditions implied by any Federal and State legislation or law of a Territory.

**APVMA Approval No: 53038/10/0804**

**Batch No:**

**DOM**

**Bar Code**

Main panel 20L bottle

---

**CAUTION**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

# Rentokil IGR Grain Protectant

ACTIVE CONSTITUENT: 30g/L S-METHOPRENE  
SOLVENT: 720g/L LIQUID HYDROCARBONS

GROUP	<b>7A</b>	INSECTICIDE
-------	-----------	-------------

For mixing with uninfested cereal grain for protection against immature stages of Lesser Grain Borer, Rust-Red Flour Beetle and Sawtoothed Grain Beetle, including organophosphate-resistant strains.

**Contents 20 Litres**

**RENTOKIL INITIAL Pty Ltd** A.C.N. 000 034 597  
150 Mowbray Road, Willoughby, NSW 2068 Australia  
a Rentokil Initial company  
Telephone 02 9370 9300

**DIRECTIONS FOR USE:**

**Restrains:** DO NOT move treated grain within 24 hours after treatment. Rentokil IGR is only effective against insects in the immature stages, not adults.

Situation/ Crop	Pest	Rate	Critical Comments
Stored Cereal grain	Lesser grain borer, rust-red flour beetle and sawtoothed grain beetle including those resistant to organophosphates.	1L in 50L of water, for 6 to 9 months protection.	Apply at rate of 1L diluted spray per tonne of grain.
	Stored grain insect pests (excluding <i>Sitophilus</i> spp.)	As above plus Rentokil Fenitrothion 1000 insecticide at the rates shown on the label.	

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

**IN WA FOR USE BY BULK HANDLING AUTHORITIES ONLY.**

**WITHHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED**

**GENERAL INSTRUCTIONS**

**Insecticide Resistance Warning**



For insecticide resistance management Rentokil IGR Grain Protectant is a Group 7A insecticide.

Some naturally-occurring insect biotypes resistant to Rentokil IGR Grain Protectant and other Group 7A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Rentokil IGR Grain Protectant or other Group 7A insecticides are used repeatedly. The effectiveness of Rentokil IGR Grain Protectant on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Rentokil Initial Pty Ltd accepts no liability for any losses that may result from the failure of Rentokil IGR Grain Protectant to control resistant insects.

Rentokil IGR Grain Protectant may be subject to specific resistance management strategies. For further information contact your local supplier, Rentokil Initial Pty Ltd representative or local agricultural department agronomist.

**MIXING:**

Using clean water only, dilute sufficient material for the job at hand. Diluted spray should be used within 48 hours.

**EQUIPMENT:**

The diluted spray should be applied through special grain spraying equipment. The spray nozzle(s) must be capable of delivering an even spray pattern, and the output of spray must be regulated according to the flow of the grain.

**CHECK RATE OF TREATMENT:**

The spray rate, measured in litres per hour must equal the auger or elevator uptake in tonnes per hour.

**CLEANING UP:**

On completion, wash out spraying equipment and mixing and measuring utensils with water.

## **EXPORT OF TREATED PRODUCE**

Users should note that suitable MRLs or import tolerances might not have been established in all markets for produce treated with Rentokil IGR. Before treating grain for export, please check with Rentokil Initial Pty Ltd for the latest information on MRLs and import tolerances.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT:** DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

**STORAGE AND DISPOSAL:** Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**SAFETY DIRECTIONS:** Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. Do not inhale vapour or spray mist. When preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length PVC gloves. Wash hands after use. After each day's use, wash gloves and contaminated clothing.

**FIRST AID:** If poisoning occurs, contact a Doctor or Poisons Information Centre. Phone Australia 131126. If swallowed do NOT induce vomiting. Give a glass of water.

**MSDS:** Additional information is listed in the Material Safety Data Sheet.

**MANUFACTURERS WARRANTY AND EXCLUSION OF LIABILITY:** This product as supplied is of high grade and believed to be suitable for any purpose for which it is expressly recommended and must be used in accordance with the Directions for Use given on the label. No responsibility is accepted in respect of the product save those non-excludable conditions implied by any Federal and State legislation or law of a Territory.

**APVMA Approval No: 53038/20/0804**

**Batch No:**

**DOM**

**Bar Code**